CS281B/Stat241B: Advanced Topics in Learning & Decision Making

Dirichlet Process Mixtures

Lecturer: Michael I. Jordan *Scribes: Wei Wang*

1 Comments on De Finetti Theorem

Theorem 1 *Exchangeability:* $p(x_1, \ldots, x_n) = p(x_{\pi(1)}, \ldots, x_{\pi(n)})$ *for any permutation* π *and for all* n*, iff*

$$p(x_1, \ldots, x_n) = \int dp(\theta) \prod_{i=1}^{n} p(x_i|\theta)$$

for some underlying distribution $p(\theta)$*, and* $p(x_i|\theta)$*.*

Notes:

- The underlying distribution $p(\theta)$ could be, for example,the Dirichlet process, $G \sim DP(\cdot)$. G is a measure on θ but also a random variable. First pick G, and then for each fixed G sample the x_i's according to $p(x_i|G)$. So an observer observing x will not know the underlying G or which sample indices correspond to which x_i's.
- Exchangeability is not equivalent to independent and identically distributed (iid). The graphical model representing exchangeability shows that $\{x_1, \ldots, x_n\}$ are not independent.



Figure 1: Exchangeability $\neq$ iid.

2 Dirichlet Process Mixtures

We have already discussed three representations of the Dirichlet Process: the stick breaking representation, the chinese restaurant process, and the urn model.



Figure 2: Dirichlet Process mixture.

- In the stick breaking representation, $G \sim DP(\alpha, G_0)$ can be written as an infinite sum of spikes $G = \sum_{i=0}^{\infty} \pi_i \delta_{\theta_i}$, where $\theta_i|\alpha, G_0 \sim G_0$, $\pi_i'|\alpha, G_0 \sim Beta(1, \alpha)$, and $\pi_i = \pi_i' \prod_{l=1}^{i-1}(1 - \pi_l)$. The π_i's can be represented as the remaining length of a stick after $i - 1$ pieces have been broken off each with size according to the Beta distribution. Here G_0 is a measure on θ_i and α determines how closely the histogram of spikes represents G_0.
- In the chinese restaurant process, x_1 starts a new table and picks (μ, σ^2) from the prior, and each subsequent x_i joins a table if it is close to (μ, σ^2), or else starts a new table with a new (μ, σ^2) chosen from the prior. Here G_0 is a measure on (μ, σ^2).
- In the urn model,

$$x_i|\theta_i \sim F(\theta_i)$$
$$\theta_i|G \sim G$$
$$G \sim DP(\alpha, G_0)$$
$$\theta_i|\theta_1, \ldots, \theta_{i-1} \sim \frac{\alpha}{i - 1 + \alpha} G_0 + \frac{1}{i - 1 + \alpha} \sum_{j=1}^{i-1} \delta(\theta_j)$$

We now introduce a fourth representation of Dirichlet Processes as the infinite limit of a finite mixture (see the Radford Neal paper).

-

$$x_i|c, \phi \sim F(\phi_{c_i})$$
$$c_i|p \sim Multinomial(p_1, \ldots, p_k)$$
$$\phi_c \sim G_0$$
$$p_1, \ldots, p_k \sim Dirichlet(\frac{\alpha}{k}, \ldots, \frac{\alpha}{k})$$



Figure 3: Finite mixture.